

DOI: https://dx.doi.org/10.51227/ojafr.2025.29

DIETARY LEUCAENA LEAVES IMPROVE GROWTH PERFORMANCE AND CARCASS QUALITY OF VIETNAMESE GOATS

Tran Ngoc LIEM¹, Ngo Mau DUNG¹, Le Minh DUC¹, Duong Thanh HAI¹, Vo Thi Minh TAM¹, Le Van AN¹, Le Thi Ouvnh ANH², Songyos CHOTCHUTIMA³, and Phoompong BOONSAEN³

¹University of Agriculture and Forestry, Hue University, 102 Phung Hung, Phu Xuan, Hue, Vietnam

²University of Economics, Hue University, 99 Ho Dac Di, An Cuu, Hue, Vietnam

³Kasetsart University, 50 Ngamwongwan Rd, Chatuchak, Bangkok 10900, Thailand

™Email: levanan@hueuni.edu.vn

Supporting Information

ABSTRACT: The experiment was conducted at a research farm for sixteen male goats, with an average body weight of 12.32 ± 0.14 kg. They were randomly allocated into 4 groups corresponding to 4 diets and fed individually. The diets were formulated to consist of 90% of Guinea grass (*Panicum maximum*) and 10% of concentrated feed as basal (in DM). Leucaena leaves were substituted at 0%, 10%, 20% and 30% of Guinea grass in four respective diets. A 2-week adaptation period was provided for the goats to the diets and feeding system before data collection. Feed intake, weight gain, feed conversion ratio, and carcass traits of goats differed significantly among the four diets (P < 0.05). The inclusion of leucaena leaves in the diets increased feed intake. As the levels of leucaena leaves in diets increased up to 30%, there were corresponding improvements in weight gain. Daily weight gain increased from 45 to 61 g/day and feed conversion ratio (FCR) decreased from 8.43 to 6.62 kg feed/kg gain. Higher leucaena inclusion improved carcass traits but did not affect loin meat quality. Economic analysis also indicated that including up to 30% leucaena leaves in the goats' diet provides a profitable outcome for farmers. The economic impact increased with the rising levels of leucaena leaves in the goats' diet. It is recommended that leucaena leaves be utilized for goat raising in smallholder farming systems in Vietnam.

Check for updates

PII: \$222877012500029-15
Received: June 26, 2025
Revised: September 12, 2025
Accepted: September 14, 2025

Keywords: Carcass, Feed conversion ratio, Goats, Growth, Leucaena.

INTRODUCTION

Goat production is an important contributor to global meat and dairy supply, particularly in developing countries where smallholders dominate (Hegde, 2020). Goats are valued for their low investment requirements, adaptability to harsh climates, and growing consumer demand for meat. However, productivity remains limited by feed shortages, protein deficiencies, and the decline of natural grazing lands (Mazinani and Rude, 2020; Lohani and Bhandari, 2021; Nguyen et al., 2023). In Vietnam, the goat population has more than doubled in the past decade, with over 417,000 households engaged in small-scale farming, yet feed scarcity continues to restrict production efficiency (Nguyen et al., 2023).

Leucaena (Leucaena leucocephala) is a perennial legume that offers a promising solution to these constraints. Rich in crude protein (20–30% DM), it produces over 6 tons DM/ha annually, adapts well to tropical environments, and can be harvested year-round with minimal inputs (Casanova-Lugo et al., 2014). Studies have demonstrated that leucaena improves feed intake, growth, and carcass yield in ruminants (Muinga et al., 1995; Wiyabot, 2022; Marhaeniyanto et al., 2023). Although it contains anti-nutritional compounds such as mimosine, these can be managed when inclusion rates are controlled (De Angelis et al., 2021).

This study aims to investigate the effects of different levels of leucaena leaves in grass-based diets on growth performance, meat quality and economic analysis of goat production in small-scale farming in Vietnam. Co goat production is primarily managed by smallholders, who feed their goats mainly by natural grasses, with leguminous forages rarely included in their diets (Nguyen et al., 2023; Olmo et al., 2024). Therefore, the objective of our experiment is to identify the optimal inclusion rates of leucaena in goat feeds that do not adversely affect animal growth.

MATERIALS AND METHODS

Animals

The experiment was conducted on 16 male goats of the local breed Co goat, at the age of 9 months, with an initial average weight of 12.32 ± 0.14 kg per goat. The goats were vaccinated against pasteurella, cholera, and foot-and-mouth disease. They were uniformly dewormed.

Diets and feeding

Guinea grass (*Panicum maximum* cv. Mombasa) and leucaena (*Leucaena leucocephala* cv. Taramba) were grown at the Institute for Development Studies, University of Agriculture and Forestry, Hue University. Guinea grass was harvested at a cutting interval of 45 days, and leucaena leaves were harvested at a cutting interval of 4 months. Only edible parts of guinea grass and leucaena leaves were collected daily, chopped, and thoroughly mixed before being fed to the goats as fresh matter. The concentrated feed was BEEF622 from the feed market. The chemical composition of the ingredients is presented in Table 1. Goats were randomly divided into 4 groups and fed according to 4 experimental diets with the levels of leucaena at 0%, 10%, 20%, and 30% on a dry matter basis (Table 2), including: Diet 1 (KP1) consists of 90% Guinea grass and 10% concentrated feed; Diet 2 (KP2) consists of 80% Guinea grass, 10% concentrated feed and 10% leucaena; Diet 3 (KP3) consists of 70% Guinea grass, 10% concentrated feed and 20% leucaena; Diet 4 (KP4) consists of 60% Guinea grass, 10% concentrated feed and 30% leucaena.

Goats were kept individually in separate pen cages, equipped with a water supply and free access to mineral blocks. They were fed twice daily at 9:00 a.m. and 3:00 p.m. The feeding process began with concentrates, which were given separately, followed by mixed green feed. The amount of feed in DM provided to goats per day was calculated at 4% of their body weight. At the end of each day, any leftover feed was collected, dried, and weighed. Every month, the amount of feed supply was adjusted according to each goat's body weight to ensure that the feed supply met their nutritional requirements. An adaptation period of 2 weeks before the feeding experiment, which followed, lasted for 4 months, from August to December 2024.

Table 1 - Chemical composition of the ingredients								
Ingredients	DM (0/)	Chemical composition (% in DM basic)						
	DM (%)	CP	NDF	ADF	Ash			
Guinea grass	24.3	8.6	73.0	40.6	8.0			
Leucaena leaves	33.2	25.4	33.0	19.2	7.0			
Concentrated feed	86.4	19.0	37.6	20.2	10.0			
DM is the abbreviation of dry m	natter CP is the abbreviat	ion for crude protein	NDE is the abbreviation	n of neutral detergent	fibre and ADE is the			

DM is the abbreviation of dry matter, CP is the abbreviation for crude protein, NDF is the abbreviation of neutral detergent fibre, and ADF is the abbreviation of acid detergent fibre.

Table 2 - Ingredient and chemical composition of diets								
Ingredient composition (kg/100 kg DM)	Experimental diets	KP1	KP2	KP3	KP4			
Concentrated feed		10	10	10	10			
Guinea grasses		90	80	70	60			
Leucaena leaves		0	10	20	30			
Total in ration		100	100	100	100			
Chemical composition (g/kg DM)*								
OM		931	930	929	929			
CP		96	113	130	146			
EE		19	22	26	29			
NDF		694	654	614	574			
ADF		386	364	342	321			
ME (MJ/kg DM)		9.34	9.61	9.88	10.15			

*Values calculated based on the composition of ingredients. OM is the abbreviation of organic matter, EE is the abbreviation of ether extract, ME is the abbreviation of metabolisable energy. Diet 1 (KP1) consists of 90% Guinea grass and 10% concentrated feed; Diet 2 (KP2) consists of 80% Guinea grass, 10% concentrated feed and 10% leucaena; Diet 3 (KP3) consists of 70% Guinea grass, 10% concentrated feed and 20% leucaena; Diet 4 (KP4) consists of 60% Guinea grass, 10% concentrated feed and 30% leucaena.

Experiment design

The experiment was conducted at the research farm of the Institute for Development Studies of the University of Agriculture and Forestry, Hue University, Vietnam, from August 2024 to January 2025. Sixteen (16) goats were randomly divided into 4 groups corresponding to 4 diets (Completely Randomized Design), raised and fed individually in 16 pens. Each pen measured 1.0 m in height, 1.5 m in length, and 0.8 m in width, and was located 0.8 m above the ground. The pens were identical in size and environmental conditions. Each pen had a separate feeder and tap water. All pens were set in an animal facility that maintained uniform environmental conditions.

Data collection

Feed intake was recorded daily for each goat. The body live weight of each goat was measured at 8:00 a.m. before feeding on days 0, 30, 60, 90, and 120 of the experiment. The prices of forages were calculated based on the actual price

of 700 VND/kg of Guinea grass and 1,000 VND/kg of leucaena leaves on the fresh material (Conversion rate: 1 USD \approx 25,000 VND in December 2024). The price of concentrated feed was determined based on the market price. After 4 months, at the end of the experiment, 12 goats were slaughtered to measure the characteristics of the carcass and the chemical composition of the loin. The measurement of carcass traits in goats was conducted following the Vietnam National Standard on Animal Welfare – Slaughter (TCVN 13905-1:2023). Before slaughter, the goats were fasted for 18 hours, provided free access to water, and measures were taken to minimize stress. The pre-slaughter weight was recorded using the Nhon Hoa Scale 30 kg (model CDH-30) with an error margin of \pm 50 g to \pm 150 g. The goats were electronically stunned before having their jugular vein and carotid artery cut. Blood was drained into a pre-weighed bucket and weighed. Hair was removed and weighed. The internal organs (digestive tract, lungs, trachea, heart, liver, kidneys, kidney fat, spleen and pancreas) were removed and weighed by Nhon Hoa Scale 10 kg (model PDM 036-2017) with an error margin of \pm 5 g to \pm 15 g. The empty body weight (excluding blood, hair, and internal organs) was measured. After the skin was removed, the head was separated at the atlas vertebra, and the legs were separated at the carpal and tarsal joints, then weighed. The hot carcass weight was measured without blood, hair, internal organs, head, legs and skin. Finally, loin meat samples were analyzed for dry matter (DM), crude protein (CP), ether extract (EE), and total ash content.

Chemical analysis

Chemical analysis of the samples was conducted to measure dry matter (DM), crude protein (CP), ether extract (EE), and total ash according to AOAC (1990). Additionally, neutral detergent fibre (NDF) and acid detergent fibre (ADF) were analyzed using the method described by Van Soest et al. (1991).

Data analysis

Collected data were statistically analyzed using ANOVA with Minitab version 16.2.0 (2010). Comparison of significant differences in mean values was assessed at the probability level of P<0.05. The statistical model used is as follows:

$$Y_{ii} = \mu + T_i + e_{ii}$$

where: μ represents the overall mean value; T_i denotes the effect of the diet; and e_{ij} is the error term.

RESULTS AND DISCUSSION

Feed intake

The daily feed intake of goats in each month and the average of 4 months differed between the diets (P<0.05; Table 3). The lowest feed intake was recorded at 385 g/day for the KP1 without leucaena, while the highest was 427 g/day for the KP2 containing 10% of leucaena in the diet (on a DM basis). The trend of feed intake decreased when leucaena leaf increased to 20 and 30%, although it remained higher than the intake for KP1 without leucaena. Previous studies have shown that substituting leucaena in diets based on grass or maize can lead to increased feed intake (Balogun and Otchere, 1995; Haque et al., 1997; Fasae et al., 2011). Wiyabot (2022) found no significant difference in feed intake when goats were fed diets with leucaena levels at 25%, 50%, 75%, and 100%. However, a diet with 40% leucaena leaves resulted in slight hair loss (Balogun and Otchere, 1995). The leucaena contents of toxicity may contribute to reduced feed intake and productivity (Phaikaew et al., 2012). In our experiment, increasing the level of leucaena leaves in the diets led to higher feed intake compared to goats fed only on guinea grass. A grass-leucaena mix increased feed intake, but excessive leucaena may reduce it (Table 3).

Table 3 - Daily feed intake (g D	M/head/day) by ı	month of experim	ent			
Daily food intake by month	Experiment diet (Mean ± Standard Deviation)					Dyelue
Daily feed intake by month	KP1	KP2	KP3	KP4	SEM	P value
First month	325 ± 22.8b	402 ± 33.1ª	366 ± 23.5ab	356 ± 20.9ab	12.8	0.009
Second month	362 ± 17.4	393 ± 30.0	379 ± 5.1	373 ± 11.8	9.25	0.191
Third month	384 ± 24.8	418 ± 22.0	417 ± 3.1	414 ± 10.5	8.73	0.054
Fourth month	433 ± 12.1	453 ± 20.8	460 ± 6.1	458 ± 14.6	7.20	0.080
Average	385 ± 18.4b	427 ± 23.4a	415 ± 7.4ab	410 ± 13.9ab	8.42	0.028
Daily feed intake of ingredients						
Concentrated feed	85.28 ± 3.22	89.47 ± 5.94	91.28 ± 1.58	88.36 ± 5.84	2.27	0.341
Guinea grass	297 ± 18.16a	283 ± 18.58a	239 ± 5.88b	198 ± 8.70°	7.01	0.001
Leucaena leaves	Oq	42.8 ± 2.82c	81.4 ± 2.00b	115.8 ± 5.08a	1.54	0.001

a, b, c values within a row with different letters were significantly different (P<0.05). Diet 1 (KP1) consists of 90% Guinea grass and 10% concentrated feed; Diet 2 (KP2) consists of 80% Guinea grass, 10% concentrated feed and 10% leucaena; Diet 3 (KP3) consists of 70% Guinea grass, 10% concentrated feed and 20% leucaena; Diet 4 (KP4) consists of 60% Guinea grass, 10% concentrated feed and 30% leucaena.

Effects of dietary levels of leucaena on the growth of goats

Increasing the level of leucaena leaves in diets that include Guinea grass and concentrated feed resulted in improved live weight and daily weight gain, while also decreasing the feed conversion ratio (FCR) (Table 4). In the first two months, there was no significant difference in the live weight of goats across four diets. However, differences in growth performance became noticeable during the third and fourth months (P<0.05). Substituting leucaena leaves into the diets led to a positive increase in the body live weight of the goats (P<0.05). During the first month of the experiment, there was no difference in daily weight gain among the diets. However, from the second month to the fourth month, differences in daily weight gain of goats emerged between the control group (KP1), which had no leucaena leaves, and the diets that included leucaena leaves (KP2, KP3, and KP4) (P<0.05). Increasing the level of leucaena leaves in the diet was associated with higher daily weight gain in the goats, with the highest gain observed at 30% leucaena leaves in KP3. The FCR improved from KP1 to KP4, decreasing from 8.43 to 6.62, respectively.

The inclusion of leucaena leaves in goat rations has been studied in many countries. Our experiment focused on local breeds "Co goat", which typically have smaller body weight and daily weight gains compared to breeds such as Back Thao, Boar, or other crossbreeds (Pham and Nguyen, 2015; Pham et al., 2019). We found that incorporating leucaena leaves at levels up to 30% in the diets of these local breeds of goat in Vietnam improved both feed intake and daily weight gain. These findings align with those of Adejumo and Ademosun (1991); Fasae et al. (2011); Marhaeniyanto et al. (2023), who also reported positive effects from including various levels of leucaena in diets. Specifically, Marhaeniyanto et al. (2023) observed that supplementing with 20% Leucaena leucocephala leaves in a concentrate containing 15% crude protein resulted in an average daily gain of 99.29 g/head/day. Conversely, Wiyabot (2022) reported no significant difference in daily weight gain among goats when leucaena was used as a roughage substitute at levels exceeding 50%. Additionally, Adejumo and Ademosun (1991) recommended that to promote growth without adverse effects, leucaena should not comprise more than 40% in goat rations. We recommend including 30% of leucaena in the diet to ensure that toxicity thresholds remain manageable for smallholder farming systems and align well with the needs of our goat breed. Additionally, biomass production of leucaena at the smallholder scale in Vietnam is limited. Therefore, our findings suggest that maintaining this 30% leucaena in the diet will optimize and sustain feed supply effectively.

e welght (kg/head)	KP1	KP2	KP3	KP4	SEM	Pvalu
Initial body weight	12.15	12.37	12.55	12.20	0.300	0.77
First month	13.05	13.45	13.68	13.42	0.294	0.530
Second month	14.20	14.90	15.22	15.08	0.304	0.138
Third month	15.95b	16.98ab	17.40a	17.40a	0.268	0.00
Fourth month	17.52b	18.73a	19.33ª	19.48a	0.258	0.00
ly weight gain (g/day)						
First month	30	36	38	41	2.98	0.12
Second month	38 ^b	48 ^{ab}	52 ^a	55ª	2.89	0.009
Third month	58 ^b	69 ^a	73ª	78 ª	2.51	0.003
Fourth month	52 ^c	58bc	64 ab	69ª	2.00	0.003
Average	45c	53b	56 ab	61 ª	1.59	0.003
FCR (kg feed/kg gain)	8.43a	7.88ab	7.20bc	6.62°	0.246	0.001

a. b. c values within a row with different letters were significantly different (P<0.05). Diet 1 (KP1) consists of 90% Guinea grass and 10% concentrated feed; Diet 2 (KP2) consists of 80% Guinea grass, 10% concentrated feed and 10% leucaena; Diet 3 (KP3) consists of 70% Guinea grass, 10% concentrated feed and 20% leucaena; Diet 4 (KP4) consists of 60% Guinea grass, 10% concentrated feed and 30% leucaena.

Carcass quality of goats

There were differences in slaughter body weight, hot carcass weight, and percentage of carcass among goats on different diets (Table 5). Increasing the level of leucaena leaves in their diets resulted in varying growth performance in the goats. The hot carcass weight was found to be higher in the rations containing 20% and 30% leucaena leaves, while the lowest hot carcass weight was observed in the rations without leucaena inclusion (P<0.05) (Table 5). The carcass weight, body meat, and their ratios in KP3 and KP4 were higher than those in KP1. Increasing the level of leucaena leaves in the diet markedly improved the carcass characteristics of slaughter goats. However, there were no differences in the meat quality of the loin based on DM, CP, EE, and total ash among diet treatments. The study concluded that incorporating leucaena leaves up to 30% in a basal goat diet of guinea grass and concentrated feed improved meat production, but had no effects on the loin meat quality in local goats. Leucaena inclusion improved carcass traits compared to traditional local goat diets in Vietnam (Pham et al., 2019). This finding aligns with the carcass characteristics observed in Afar goats fed with leucaena in Ethiopia (Terefe et al., 2013; Gebrehiwot et al., 2017) and in indigenous Anglo-Nubian hybrid goats in Thailand (Wiyabot, 2022).

Economic analysis

leucaena; Diet 4 (KP4) consists of 60% Guinea grass, 10% concentrated feed and 30% leucaena.

The cost-benefit analysis of investing in this experiment was calculated based on the prices of animals, feed, and veterinary services for goats during their feeding period. Table 6 provides a summary of the economic impact data on sixteen male goats raised on four different rations. The lowest economic impact was found in KP1. As the inclusion of leucaena leaves in the diets increased from 10%, 20%, to 30%, the economic analysis results increased by 28%, 43% and 62%, respectively, compared to the baseline without leucaena leaves. In the condition of market price fluctuations, we conducted a sensitivity analysis to evaluate how the profit changed in response to a 10% increase or 10% decrease in the sale price. Our findings indicate that, in all scenarios, KP2, KP3, and KP4 consistently generate higher profits than KP1 (Table 6). The cost-benefit analysis of goat farming in this experiment indicates that goat farming can be a profitable business for smallholders in Vietnam when incorporating leucaena leaves at up to 30% dry matter (DM) in their forage. Leucaena is primarily fed to goats as fresh material through cut-and-carry feeding systems, which are both flexible and laborefficient (Palmer et al., 2010). Under tropical conditions, leucaena can provide over 4 tonnes of foliage per hectare per year (Casanova-Lugo et al., 2014; Cowley and Roschinsky, 2019), making it a valuable feed source for small-scale goat production.

Experimental diets Category	KP1	KP2	КРЗ	KP4	SEM	P value
Pre-slaughter weight (kg)	17.53b	18.57ab	19.20a	19.60a	0.324	0.009
Blood (kg)	0.60	0.61	0.61	0.61	0.027	0.992
Hair (kg)	0.03	0.05	0.04	0.04	0.006	0.122
Internal organs (kg)	5.80	5.45	5.54	5.41	0.092	0.059
Empty body weight (kg)	10.54°	11.48bc	12.22ab	12.89ª	0.279	0.002
Head (kg)	1.28b	1.43 ab	1.48a	1.54a	0.036	0.006
Legs (kg)	0.50	0.51	0.53	0.60	0.026	0.102
Skin (kg)	1.29b	1 .38 ^{ab}	1.54ab	1.62a	0.062	0.021
Hot carcass weight (kg)	7.47°	8.15bc	8.66ab	9.11a	0.205	0.003
Dressing percentage (%)						
Pre-slaughter weight bases (%)	42,59°	43.88bc	45.12ab	46.45a	0.529	0.005
Empty body weight bases (%)	70.85	71.02	70.91	70.76	0.470	0.982
Chemical composition of loin						
DM (%)	22.55	23.19	23.27	23.23	1.74	0.989
CP (%)	84.72	85.31	86.31	85.70	0.587	0.388
EE (%)	3.78	4.35	3.05	4.02	0.434	0.320
Total ash (%)	4.64	4.59	4.73	5.58	0.210	0.951

Table 6 - Economic analysis of goat raising by diets **Experimental diets Price** Unit (VND)* KP1 KP2 KP3 KP4 Investment costs Animal breed 120,000 1,458,000 1,485,000 1,506,000 1,464,000 kg 142,392 Concentrated feed kg 13,000 133,032 139,571 137,846 Guinea grass kg 700 113,466 107.872 91,164 75,647 Leucaena leaves 1,000 17,123 32,559 46,315 kg Vaccine 50,000 50,000 50,000 50,000 50,000 head **Total** 1,754,498 1,799,566 1,822,115 1,773,808 Sale income Fixed sale price 120,000 kg VND 2.103.000 2.247.000 2.319.000 2.337.000 Income **Profit VND** 348.502 447.435 496.885 563.192 % compared to KP1 100 128 143 162 Sensitivity analysis Sale price increases by 10% 132.000 kg Income **VND** 2,313,000 2,472,000 2,551,000 2,571,000 Balance/Profit 672,434 **VND** 558.502 728.885 797,192 % compared to KP1 100 120 131 143 Sale price decreases by 10% kg 108,000 VND 1,893,000 2.022.000 2,087,000 2,103,000 Income Balance/Profit **VND** 138,502 222,434 264,885 329,192 % compared to KP1

Conversion rate: 1 USD ≈ 25,000 VND in December 2024. Diet 1 (KP1) consists of 90% Guinea grass and 10% concentrated feed; Diet 2 (KP2) consists of 80% Guinea ss, 10% concentrated feed and 10% leucaena; Diet 3 (KP3) consists of 70% Guinea grass, 10% concentrated feed and 20% leucaena; Diet 4 (KP4) consists of 60% Guinea grass, 10% concentrated feed and 30% leucaena

CONCLUSIONS

This study demonstrated that feed intake, weight gain, feed conversion ratio, and carcass traits of goats differed significantly among the four dietary treatments. The inclusion of leucaena leaves in the diets increased feed intake and improved growth performance as the proportion of leucaena rose to 30%. Daily weight gain increased from 45 g/day in the control diet to 61 g/day in the KP4 diet, while the feed conversion ratio improved from 8.43 to 6.62 kg feed/kg gain. Higher levels of leucaena also enhanced carcass traits, although loin meat quality remained unaffected. Economic analysis confirmed that supplementing up to 30% leucaena leaves in goat diets yields profitable outcomes for smallholder farmers, with greater economic benefits at higher inclusion levels. Overall, the findings support the recommendation that leucaena leaves be incorporated into goat feeding strategies to improve productivity and profitability in smallholder farming systems in Vietnam.

DECLARATIONS

Corresponding author

Correspondence and requests for materials should be addressed to Le Van AN; Email: levanan@hueuni.edu.vn; ORCID: https://orcid.org/0000-0002-0954-6208.

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

TN.Liem, NM.Dung, and LM.Duc were responsible for implementing the research activities. DT.Hai and VTM.Tam carried out the laboratory work, while LTQ.Anh conducted the economic analysis. S.Chotchutima and P.Boonsaen collaborated as partners in developing the research plan and provided funding through the project "Improving Smallholder Goat Fattening Systems Based on Fodder from Grasses and Legumes in Thailand, Laos, and Vietnam", supported by the Lancang-Mekong Cooperation Special Fund.

Ethical regulations

All experimental procedures involving animals were conducted in accordance with Vietnamese regulations on animal welfare and research ethics and were approved by the Animal Ethics Committee of Hue University (Approval No. HUVN0055, dated 20 February 2025). The authors also complied with the ARRIVE guidelines.

Funding

This study was funded by the "Lancang-Mekong Cooperation Special Fund" and the research fund of Hue University, code DHH2024-02-180.

Acknowledgements

The authors express thankfulness for the cooperation program on "Improving smallholder goat fattening systems based on fodder from grasses and legumes in Thailand, Laos and Vietnam" funded by the "Lancang-Mekong Cooperation Special Fund" and the research fund of Hue University, code DHH2024-02-180.

Competing interests

The authors declare no competing interests in this research and publication.

REFERENCES

Adejumo JO and Ademosun AA (1991). Utilization of leucaena as supplement for growing dwarf sheep and goats in the humid zone of west Africa. Small Ruminant Research, 5(1-2):75-82. https://doi.org/10.1016/0921-4488(91)90032-L

AOAC (1990). Official Method of Analysis. 15th Edition. Association of Official Analytical Chemists, Washington DC.

Balogun RO, and Otchere, EO (1995). Effect of level of Leucaena leucocephala in the diet on feed intake, growth and feed efficiency of Yankasa rams. Tropical Grasslands, 29: 150-154. https://www.tropicalgrasslands.info/public/journals/4/Historic/Tropical%20Grasslands%20Journal%20archive/PDFs/Vol_29_1995/Vol_29_03_95_pp150_154.pdf

Casanova-Lugo F, Petit-Aldana J, Solorio-Sánchez FJ, Parsons D and Ramírez-Avilés L (2014). Forage yield and quality of *Leucaena leucocephala* and *Guazuma ulmifolia* in mixed and pure fodder banks systems in Yucatan, Mexico. Agroforestry Systems, 88: 29-39. https://doi.org/10.1007/s10457-013-9652-7

Cowley FC and Roschinsky R (2019). Incorporating leucaena into goat production systems. Tropical Grasslands-Forrajes Tropicales, 7(2): 173–181. https://doi.org/10.17138/tgft(7)173-181

De Angelis A, Gasco L, Parisi G and Danieli PP (2021). A multipurpose leguminous plant for the Mediterranean countries: Leucaena leucocephala as an alternative protein source: A review. Animals, 11(8): 2230. https://doi.org/10.3390/ani11082230

- Fasae OA, Adesope AI and Ojo VOA (2011). The effect of Leucaena leaf supplementation to maize residues on village goat performance. Journal of Animal & Plant Sciences, 10(2): 1276- 1282. https://www.m.elewa.org/JAPS/2011/10.2/2.pdf
- Gebrehiwot G, Negesse T and Abebe A (2017). Effect of feeding Leucaena leucocephala leaves and pods on feed intake, digestibility, body weight change and carcass characteristic of central-highland sheep fed basal diet wheat bran and natural pasture hay in Tigray, Ethiopia. International Journal of Agriculture, Environment and Biotechnology, 10(3): 367-376. https://ndpublisher.in/admin/issues/IJAEBv10n3n.pdf
- Haque N, Khan MY and Murarilal (1997). Effect of level of Leucaena leucocephala in the diets of jamunapari goats on carbon nitrogen and energy balances. Asian-Australasian Journal of Animal Sciences, 10(5): 455-459. https://doi.org/10.5713/ajas.1997.455
- Hegde NG (2020). Goat development: an opportunity to strengthen rural economy in Asia and Africa. Asian Journal of Research in Animal and Veterinary Sciences, 5(4): 30–47. https://ssrn.com/abstract=4348817
- Lohani M and Bhandari D (2021). The importance of goats in the world. Professional Agricultural Workers Journal (PAWJ), 6(2): 9-21. DOI: https://tuspubs.tuskegee.edu/pawj/vol6/iss2/4
- Marhaeniyanto E, Susanti S and Hidayati A (2023). Using different-level of *Leucaena leucocephala* leaves in concentrated feeds to increase goat farming production. Jurnal Ilmu- Ilmu Peternakan (Indonesian Journal of Animal Science), 33(2): 178-187. DOI: https://doi.org/10.21776/ub.jiip.2023.033.02.05
- Mazinani M and Rude B (2020). Population, world production and quality of sheep and goat products. American Journal of Animal and Veterinary Sciences, 15(4): 291-299. https://doi.org/10.3844/ajavsp.2020.291.299
- Muinga RW, Topps JH, Rooke JA and Thorpe W (1995). The effect of supplementation with Leucaena leucocephala and maize bran on voluntary food intake, digestibility, live weight and milk yield of Bos indicus × Bos taurus dairy cows and rumen fermentation in steers offered Pennisetum purpureum ad libitum in the semi-humid tropics. Animal Science, 60(1): 13-23. https://doi.org/10.1017/S1357729800008080
- Nguyen VD, Nguyen CO, Chau TML, Nguyen DQD, Han AT and Le TTH (2023). Goat production, supply chains, challenges, and opportunities for development in Vietnam: A Review. Animals, 13(15): 2546. https://doi.org/10.3390/ani13152546
- Olmo L, Nguyen HV, Nguyen XB, Bui TN, Ngo CTK, et al. (2024). Goat meat supply and demand in Vietnam: global context and opportunities and risks for smallholder producers. Animal Production Science, 64: AN23416. https://doi.org/10.1071/AN23416
- Palmer B, Jones RJ, Poathong S and Chobtang J (2010). The value of *Leucaena leucocephala* bark in leucaena-grass hay diets for Thai goats. Tropical Animal Health and Production, 42: 1731–1735. https://doi.org/10.1007/s11250-010-9628-9
- Phaikaew C, Suksaran W, Ted-arsen J, Nakamanee G, Saichuer A, Seejundee S, et al. (2012). Incidence of subclinical toxicity in goats and dairy cows consuming leucaena (Leucaena leucocephala) in Thailand. Animal Production Science, 52(4): 283-286. https://doi.org/10.1071/AN11239
- Pham KD and Nguyen BM (2015). Physical appearance and growth performance of indigenous goat Co, F1 (Bach Thao×Co) and three way crossbred Goat [Boer×(Bach Thao×Co)] raised in Nho Quan, Ninh Binh Province. Journal of Science & Development, 13(4): 551-559. https://tapchi.vnua.edu.vn/wp-content/uploads/old/2472015-TC%20so4.201509cn.pdf
- Pham TH, Le AD, Tran QH and Tran QH (2019). Yield and quality of meat of Co, Bach Thao and F1 (Bach Thao X Co) goat raised in Dak Lak. Advances in Ecological and Environmental Research, 4(8): 231-240. https://www.ss-pub.org/wp-content/uploads/2019/08/AEER2019031901.pdf
- Terefe E, Yaqob Y, Dessalegn K, Tafa A, Kifle A, Gebregziabher W, et al. (2013). Market weight and carcass characteristics of intact yearling afar goats under semi-intensive feeding management. International Journal of Livestock Production, 4(6): 95-101. https://doi.org/10.5897/IJLP12.023
- Van Soest PJ, Robertson JB and Lewis BA (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. Journal of dairy science, 74(10): 3583-3597. https://doi.org/10.3168/jds.S0022-0302(91)78551-2
- Wiyabot T (2022). Management and value-added of goat production, Thailand: The Leucaena (Leucaena leucocephala) as roughage source on performance and meat quality in rainy season. Iranian Journal of Applied Animal Science, 12(4): 753-759. https://journals.iau.ir/article_697777.html

Publisher's note: Scienceline Publication Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit https://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2025